

Historic, Archive Document

Do not assume content reflects current scientific knowledge, policies, or practices.

U.S. Department
of Agriculture

APHIS-PPQ

APHIS 81-50

September 1987

PESTS NOT KNOWN TO OCCUR IN THE UNITED STATES OR OF LIMITED
DISTRIBUTION NO. 88: PLUM POX VIRUS

Prepared by L. W.H. Chang, Technical Information Systems Staff,
PPQ, APHIS, USDA, Federal Building Room 611, Hyattsville, MD
20782

Disease

Plum pox, pox, sharka, or sarka of apricot, peach, Japanese
plum, or plum

Pathogen

Plum pox virus (PPV) Atanasoff 1932

Other Names

Prunus virus 7 Christoff 1938

Group

Potyvirus. Cryptogram R/1:3.5/6.2+0.4:E/E:S/Ap (Nemeth 1986)

Economic
Importance

Plum pox virus causes serious losses to plums, apricots,
peaches, and Japanese plums in Europe. Fruit loss can be total
in susceptible cultivars. PPV reduces the number of fruit by
affecting pollination and increasing fruit drop. Average plum
yield per tree dropped about 83 percent from 1949 to 1963 in a
heavily infested district of Czechoslovakia. The fruit was
unsalable (Blattny and Heger 1965). External and internal
appearance, flavor, and shelflife decline (Pemberton 1980,
Zawadzka and Millikan 1971).

As there is no cure or treatment, infected trees are usually
removed. The costs of replacing diseased trees with healthy
ones, of production loss due to replacement, and of intensive
inspection and survey must also be considered part of the
economic losses. Tree replacement in one orchard in Poland for
a very susceptible plum cultivar totaled about 18 percent in 15
years (Grzyb 1984). Replacement is much higher in the Balkans,
where the disease is more severe (P. R. Fridlund, pers. comm.).

Hosts

Woody hosts appear to be restricted to the genus Prunus.
Species reported as naturally infected include P. armeniaca (= P. divaricata (European and Mediterranean Plant Protection Organization 1970), apricot (Mathys 1974)); P. cerasifera, myrobalan plum (European and Mediterranean Plant Protection Organization 1970, Mathys 1974); P. domestica, plum (Mathys 1974), P. domestica subsp. insititia, damson plum (Trifonov 1978); P. dulcis, almond (Llacer et al. 1985b); P. glandulosa (Nemeth and Schmelzer 1972); P. persica, peach (Mathys 1974); P. salicina (= P. triflora), Japanese plum (Llacer et al. 1985b); P. spinosa, blackthorn, a symptomless carrier (Jordovic et al. 1971); and P. tomentosa (European and Mediterranean Plant Protection Organization 1970).

Infected herbaceous plants may serve as a potential inoculum
source, but no natural transmission of PPV from herbaceous to
Prunus hosts has been demonstrated although experimental

transmission has occurred. Herbaceous species naturally infected in orchards include: Campanula rapunculoides, creeping bellflower (Kroll 1973); Chenopodium sp. (Llacer et al. 1985b); Lamium album, white deadnettle; Lupinus albus, white lupine; Lycium barbarum (= L. halimifolium), Barbary matrimonyvine (Kroll 1973); Dimorphotheca aurantiaca hybrida (Zawadzka and Smolarz 1978); Medicago lupulina, black medic; Melilotus officinalis, yellow sweetclover (Kroll 1973); Pisum sativum, garden pea; Ranunculus acer (Zawadzka and Smolarz 1978); Silene vulgaris, bladder campion (Kroll 1975); Solanum dulcamara, bittersweet nightshade; Trifolium incarnatum, crimson clover (Kroll 1973); T. pratense, red clover (Zawadzka and Smolarz 1978); T. repens, white clover (Kroll 1973); and Zinnia violacea (= Z. elegans), zinnia (Zawadzka and Smolarz 1978).

General
Distribution

Unless otherwise cited, the European and Mediterranean Plant Protection Organization (1983) and the Commonwealth Mycological Institute (1970) listed the following countries: Albania, Austria, Belgium (local infestation under eradication--van



Plum pox virus distribution map.

Melckebeke pers. comm. 4/15/87), Bulgaria, Cyprus (detected in 1982; eradication underway--European and Mediterranean Plant Protection Organization 1984), Czechoslovakia, Denmark (only Fyn Island, eradication underway--European and Mediterranean Plant Protection Organization 1986), East Germany, France (Commonwealth Mycological Institute 1980), Greece, Hungary, Italy, Poland, Portugal (local infestation under eradication--European and Mediterranean Plant Protection Organization 1985), Romania, Soviet Union (Crimea, Moldavia, West Ukraine--Pomazkov and Abramova 1971), Spain (Llacer et al. 1985b), Switzerland, Syria (Food and Agriculture Organization of the United Nations 1986), Turkey (Sahtiyanci 1969), United Kingdom (England and Wales--Pemberton 1980, D. L. Ebbels, pers. comm. 3/31/87), West Germany, and Yugoslavia.

Characters

Virus particles filamentous (Fig. 1), about 725-760 nm long by 20 nm wide. Three strains exist: necrotic, intermediate, and yellow (Sutic et al. 1971). Needlelike and pinwheel inclusions in nucleus and cytoplasm of leaves and ripe fruit of peach and plum. Inclusions absent from sharka-free and from pseudopox fruit (Kegler and Schade 1971, van Oosten 1971, 1972).

Transmission - Through grafts, sap inoculation (Kegler and Schade 1971) and Prunus seed. Seed transmission rates ranged up to 13.9 percent (Coman and Cociu 1976 from Nemeth 1986; Mathys 1974, Nemeth and Kolber 1983). Coman and Cociu (1976) reported 20-80 percent transmission for pollen.

(Fig. 1)



PPV particles from peach leaf cells after differential centrifugation and negative staining with phosphotungstic acid (X 33,600)(From Maroquin and Rassel 1976).

Several aphid species transmit PPV in a nonpersistent, styletborne manner (Kassanis and Sutic 1965; Kunze and Krczal 1971). No transmission to aphid progeny has been reported. All of the following vectors except four species were found by Kunze and Krczal (1971) and Krczal and Kunze (1972): Aphis citricola Van der Goot (= A. spiraecola Patch), spirea aphid (Leclant 1973); A. craccivora Koch, cowpea aphid (Leclant 1973); Brachycaudus cardui (L.), thistle aphid; B. helichrysi (Kaltenbach), leafcurl plum aphid; Hyalopterus pruni (Geoffroy), mealy plum aphid (Minoiu 1973); Myzus persicae (Sulzer), green peach aphid; M. varians Davidson (Leclant 1973); and Phorodon humuli (Schrank), hop aphid.

After 3 hours of starvation, the aphid acquires PPV (Kunze and Krczal 1971, Krczal and Kunze 1972) in 1-5 minutes of feeding (Kassanis and Sutic 1965, Leclant 1973, van Oosten 1970). PPV persists for 1 hour in M. persicae and P. humuli, but persists for 3 hours in P. humuli if the aphid is starved for 3 hours before and after acquisition (Krczal and Kunze 1972).

Testing - PPV is difficult to detect due to uneven distribution and low concentration of virus in the host. Enzyme-linked immunosorbent assay (ELISA) can detect low PPV concentrations in roots, bark, fruit, seed, flowers, and leaves. Because results are more reliable after, rather than before, foliar symptoms appear (Hamdorf 1983) and because of uneven virus distribution, ELISA is limited to confirming the presence, but not the absence, of PPV (Adams 1978a, Clark et al. 1976, Nemeth and Kolber 1983, Roggero and Lenzi 1985, Torrance 1981).

Slower tests include transmission to sensitive indicator plants (indexing). Grafting infected buds on seedlings of woody indicators (P. persica and P. tomentosa) in early spring results in symptoms in 1-8 weeks (European and Mediterranean Plant Protection Organization 1983, Nemeth 1986). The latter species served as a reliable woody indicator for 18 PPV isolates and separated PPV from 3 other viruses of plum (Rankovic 1975, 1980). Sap inoculation to the herbaceous indicator Chenopodium foetidum reliably produces symptoms only in spring in 6-8 days (European and Mediterranean Plant Protection Organization 1983).

Characteristic Damage

Infected leaves may show chlorotic flecks, bands, rings, and mottling or vein clearing. Infected fruits are discolored and drop prematurely. Symptoms vary with the locality, season, Prunus species, and cultivar. Some cultivars are symptomless.

(Fig. 2-3)



2



3

PPV symptoms on 'Canino' apricot. 2. Leaves. 3. Fruit (Courtesy J. Foster).

(Apricot)

Apricot leaves develop diffuse pale green rings and lines (Fig. 2). Fruit are deformed with chlorotic yellow rings on the skin (Fig. 3). Yellow rings and spots on the stones (Dunez 1987) separate plum pox from other apricot pox (Mathys 1974). Affected trees may be sensitive to drought and cold (Christoff 1958).

(Figs. 4-5)



4



5

Symptoms on 'Red Beaut' Japanese plum. 4. Leaves (far left leaf is healthy). 5. Fruit (From Llacer et al. 1985a).

(Japanese plum)

Japanese plum leaves may show chlorotic rings, bands, and mottling (Fig. 4), later becoming necrotic. Fruit (Fig. 5) show rings, flecks, depressed brownish areas, and conspicuous malformations at ripening (Llacer et al. 1985b).

(Figs. 6-7)



Symptoms on peach. 6. Leaves of 'GF 305'. 7. Fruit of 'Rojo del Rito' (6, courtesy J. Foster; 7 from Llacer et al. 1985a).

(Peach)

Peach leaves on young shoots show secondary and tertiary vein clearing in early spring. Leaves may be malformed (Fig. 6) with a necrotic main vein and may show some epinasty. Peach fruit develop spots and rings (Fig. 7): white or green on white-fleshed cultivars and greenish yellow to green on yellow-fleshed cultivars. Symptoms also disappear quickly (European and Mediterranean Plant Protection Organization 1983).

(Plum)

Plum leaves may show irregular vein clearings or a dark oak leaf pattern bordering the main and secondary veins. Later, characteristic yellowish to olive green flecks, bands, rings, and mottling appear (Fig. 8). Ring edges are typically distinct on the inner edge and blurred on the outer. Symptoms fade during hot months except in some cultivars where the rings and spots later develop reddish brown margins (Nemeth 1986, European and Mediterranean Plant Protection Organization 1983).


Plum fruit develop red flecks and broad rings on light-skinned fruit; flecks disappear after a short time on dark-skinned cultivars (European and Mediterranean Plant Protection

(Figs. 8-12)


8




9




10



12



11



Symptoms on plum. 8. Leaves. 9-12. Fruit. 9. Bands on 'Fleuriana'; 10. Thin, red rings on 'Early Laxton'; 11. Thin, red lines on 'Early Laxton'; 12. Grooves and pits (8 and 12, courtesy B. Zawadzka, Research Institute of Pomology and Floriculture, Skierniewice, Poland; 9-11 from van Oosten 1971b).

Organization 1983). Some cultivars with fruit that are orange, red, or purple (not green or yellow) show broad red/purple bands--diffuse on one side, distinct on the other side (Fig. 9) with thin red rings (Fig. 10) and lines (Fig. 11) near the fruit apex 2-4 weeks before picking. These symptoms are most apparent before the color darkens (van Oosten 1971, 1972).

Plum fruit may show bluish grooving and pitting (Fig. 11) with brownish-red necrosis of the underlying tissue and internal gummosis to the stone. Rings and spots may occur on the stone. Depending on the cultivar, fruit may drop prematurely (Nemeth 1986). In some cultivars, these symptoms are not always an indication of plum pox (van Oosten 1972).

To help distinguish similar symptoms caused by PPV and other viruses such as apple chlorotic leafspot virus (incitant of pseudopox), plum line pattern virus, Prunus necrotic ringspot virus, and prune dwarf virus, see tables in Kegler and Schade (1971) and Mathys (1974).

Detection Notes

Movement of infected propagative material can introduce PPV into new areas. Besides infected vegetative material (plants, roots, scions, and budwood), infected seeds of apricot, peach, and plum may pose a risk in that the resulting young trees may become inoculum sources. Domestic aphids may acquire and spread the virus from imported Prunus cut flowers to nearby fruit trees or weeds. Or, the virus may enter this country via aphid vectors hitchhiking on infected cut flowers. Infected weed hosts would be a problem if the plants were moved about.

Several problems impede early detection of infected material, increasing the probability of introduction and establishment. Roots or other parts may be infected, but the tree appears symptomless. Several years may lapse before symptoms become obvious. Careful inspection is needed to spot the earliest symptoms on newly infected trees before secondary spread occurs to uninfected trees. Symptoms may be difficult to spot because the virus spreads incompletely in the tree and may remain in a small part, for example, a twig for some time. Symptoms may be indistinct and disappear in high summer temperatures. Fruit symptoms may be apparent for less than a month before harvest (Adams 1978b, Pemberton 1980).

To minimize the risk of introducing PPV, propagative material and cut flowers of its primary Prunus hosts is prohibited or restricted entry into the United States by Title 7, Part 319.37 of the Code of Federal Regulations. Propagative material is enterable under USDA permit for scientific purposes subject to stringent entry conditions including testing for PPV.

Survey--

1. Between late spring and early summer and in fall, check young trees for leaves with chlorosis or vein clearing at least twice a season, and inspect infested orchards at least once every 3 weeks. Inspect fruiting and mother trees (for propagation of budwood) several times a year (Pemberton 1980).

2. Watch for heavy, premature fruit drop. Examine fruit during the month before harvest; also inspect dropped or harvested fruit. Look for lines, rings or diffuse mottling on fruit and rings and spots on stones.

Submit for diagnosis, suspect plant material labeled and packaged to preserve its freshness as long as possible.

Biology and Etiology

Infected Prunus trees provide the major source of PPV inoculum. From infected Prunus material, the virus is spread through grafts and budding material during propagation, or by its aphid vectors.

The number of infected trees corresponds to the number of winged vectors present on host fruit trees during the year. Beginning in late winter or early spring, overwintered aphid eggs hatch on PPV-infected fruit trees. After a few wingless generations, winged forms appear, probe or feed on the fruit tree host, and fly to other hosts to also probe and feed. In midseason, the aphids migrate to herbaceous hosts. When these plants slow their growth or die, or when the season cools in late summer, the winged generation develops and flies to stone fruit trees to lay overwintering eggs. These adults move frequently from tree to tree, probing and feeding for short periods (Kunze and Krczal 1971, Pemberton 1980, Pomazkov and Abramova 1971).

Long-distance virus dissemination by the vector may be possible. As mentioned earlier, P. humuli can transmit the virus 3 hours after acquisition if starved before and after feeding. Migrants fast for such periods in traveling to and from a distant infection source (Krczal and Kunze 1972).

Once a tree is inoculated, incubation may take 9-13 months. Systemic spread is slow and may take another 2-3 years in a young tree (Christoff 1958, European and Mediterranean Plant Protection Organization 1983).

Diseased trees first appear in a previously uninfested orchard in a random pattern. After 2 years, however, newly diseased trees appear near trees found infected in the previous year (Grzyb 1984, Jordovic 1975). This pattern seems to correspond to winged vectors introducing inoculum from outside of the orchard in the first few years, followed in later years by vector dissemination of the virus from infected trees within the orchard (Jordovic 1975).

Based on the pattern of spread to nearby trees, the estimated rate of spread averaged 10-15 m annually in the Soviet Union (Pomazkov and Abramova 1971). In Yugoslavia, the rate of infection over 10 years in 2 orchards ranged from 49 to 100 percent in orchards 100 m or less from a source of heavily infected trees, but dropped to less than 2 percent in orchards 500 m or more from the infection source. Besides distance from the source, some of the other factors affecting rate of spread are the number of infected trees (Jordovic 1968), climate, the tree cultivar, and the virus strain involved (European and Mediterranean Plant Protection Organization 1983). Spread in nurseries occurs rapidly because of such practices as planting material closely together and using undetected infected rootstocks or other infected material for propagation (Pemberton 1980).

Control

Using propagative material that is "virus-free" is an important means of exclusion. Planting orchards distant from infested orchards, planting cultivars tolerant to the virus, practicing vector control (Kegler et al. 1978), and roguing diseased trees are some of the other practices used to keep PPV under control.

Diseased trees should be eliminated before fall migrants return and disseminate PPV from these trees to healthy trees. Removing the diseased tree and surrounding symptomless trees was recommended (Kunze and Krczal 1971). All parts are best removed. Otherwise, burn leafy portions, and treat stump and suckers with systemic herbicide until no suckers emerge as suckers are usually infected. Replanting may then proceed (Adams 1978b, Pemberton 1980).

Roguing diseased trees in orchards with 10 percent infection or less kept PPV under control. In orchards with more than 10 percent infection, however, roguing was considered impractical and removing the entire orchard was recommended because that number of diseased trees meant too many trees to replace and too many latent infections that could serve as inoculum sources. That level also threatened adjacent orchards (Pemberton 1980).

Literature Cited

Adams, A. N. The detection of plum pox virus in Prunus species by enzyme-linked immunosorbent assay (ELISA). Ann. Appl. Biol. 90(2):215-221; 1978a.

_____. The incidence of plum pox virus in England and its control in orchards. Scott, P. R.; Bainbridge, A., editors. Plant disease epidemiology. Oxford, United Kingdom: Blackwell Scientific Publications; 1978b: 213-219.

- Blattny, C.; Heger, M. Some remarks to the economical importance of sarka disease in Czechoslovakia. Zast. Bilja 16(85-88):417-418; 1965 (Serbo-Croatian summary).
- Christoff, A. Die obstvirozen in Bulgarien. Phytopathol. Z. 31:381-436; 1958.
- Clark, M. F.; Adams, A. N.; Thresh, J. M.; Casper, R. The detection of plum pox and other viruses in woody plants by enzyme-linked immunosorbent assay (ELISA). Acta Hortic. 67:51-57; 1976.
- Coman, T.; Cociu, V. Transmission de la Sharka par le pollen et par les graines. Bull. Inf. Sharka. No. 2:15-21; 1976.
- Commonwealth Mycological Institute. Pathogen: Plum pox virus Atanasoff. Distribution maps of plant diseases. No. 392, 2d ed. London: Commonwealth Agricultural Bureaux; 1970. Additions and corrections; 1980.
- Dunez, J. Plum pox disease of stone fruit trees. Rome: Food and Agriculture Organization of the United Nations; 1987.
- European and Mediterranean Plant Protection Organization. Phytosanitary position in regard to the sharka disease (plum pox). Paris: EPPO Publication B, No. 66; 1970.
- _____. Data sheets on quarantine organisms. Plum pox virus. EPPO Bull. 13(1):1-7; 1983.
- _____. Plum pox virus on apricot and peach. FAO Plant Prot. Bull. 32(2):77; 1984.
- _____. Plum pox virus on apricot. Outbreaks and new records. FAO Plant Prot. Bull. 33(1):45; 1985.
- _____. Denmark: sharka virus on plum. Outbreaks and new records. FAO Plant Prot. Bull. 34(4):214; 1986.
- Food and Agriculture Organization of the United Nations. Pest and disease records. Asia and Pacific Plant Prot. Comm. Q. Newsl. 29(3):29; 1986.
- Grzyb, Z. S. Spread of the plum pox virus in the Italian prune orchard. Fruit Sci. Rep. 11(4):155-158; 1984.
- Hamdorf, G. The detection of plum pox virus (PPV) by indicator plants and enzyme-linked immunosorbent assay (ELISA). Acta Hortic. 130:151-159; 1983.

Jordovic, M. Effect of sources of infection on epidemiology of sarka (plum pox) virus disease. Deut. Akad. Landwirtschafts-swiss. Tagungsber. 97:301-308; 1968.

_____. Localization of plum orchards as a factor of the prevention of the spread of sharka virus. Jugosl. Vocarstvo 16(1/2): 89-92; 1982 (In Serbo-Croatian; English summary).

Jordovic, M.; Festic, H.; Rankovic, M. Prunus spinosa L. as host of sharka virus: isolation and some properties of the virus. Eur. Symp. Fruit Tree Virus Dis. Proc. 8th. Paris: Institut National de la Recherche Agronomique; 1971: 179-184.

Jordovic, M. M. Study of sharka spread pattern in some plum orchards. Acta Hortic. 44:147-154; 1975.

Kassanis, B.; Sutic, D. Some results of recent investigations on sarka (plum pox) virus disease. Zast. Bilja 16(85-88): 335-340; 1965 (Serbo-Croatian summary).

Kegler, H.; Schade, C. Plum pox virus. C.M.I./A.A.B. Descr. plant viruses. No. 70; 1971.

Kegler, H.; Schimanski, H.; Verderevskaja, T. D.; Trifonov, D. Zur Bedeutung der Toleranz von Obstgehölzen gegenüber virosen und Mykoplasmosen. Nachrichtenbl. Pflanzenschutz DDR 32(12): 248-251; 1978 (Russian and English summaries).

Krczal, H.; Kunze, L. Experiments on the transmission of plum pox virus by aphids. Mitt. Biol. Bundesanst. Land Forstwirtsch. Berlin-Dahlem 144:71-83; 1972 (In German; English summary).

Kroll, J. Natural and experimental host plants of the sharka virus of plum trees. Plant Virology. Proc. 7th Conf. Czechoslovak Plant Virologists. Bratislava: Slovak Academy of Sciences 1971: 397-401; 1973.

_____. Examinations concerning the range of host plants of the plum pox virus (PPV). Zentralbl. Bakteriол. Parasit. Infektiowskr Hyg. Zweite Naturwiss. Abt. 130(3):219-225; 1975 (In German; English summary).

Kunze, L.; Krczal, H. Transmission of sharka virus by aphids. Eur. Symp. Fruit Tree Virus Dis. Proc. 8th. Paris: Institut National de la Recherche Agronomique; 1971: 255-260.

Leclant, F. Ecological aspect of sharka (plum pox) virus transmission in southeastern France: additional aphid vectors. Ann. Phytopathol. 5(4):431-439; 1973 (In French; English and German summaries).

Llacer, G.; Cambra, M.; Lavina, A. Detection and preliminary studies on the sharka virus in Spain. An. Inst. Nac. Invest. Agrar. Agric. 28(3):157-166; 1985a (In Spanish; English Summary).

_____. Detection of plum pox virus in Spain. EPPO Bull. 15(3):325-329; 1985b (French and Russian summaries).

Mathys, G., editor. Progress obtained through the knowledge of sharka. EPPO Bull. 4(1); 1974 (In French).

Minoiu, N. The vectors transmitting plum pox virus (Prunus virus 7 Christ) to plum. An. Inst. Cer. Pentru. Prot. Plant 9:49-56; 1973 (In Romanian; English summary).

Nemeth, M. Virus, mycoplasma and rickettsia diseases of fruit trees. Boston: Martinus Nijhoff Publishers; 1986: 463-479. Szalay-Marzso, L., translator; translation revised by A. F. Posnette. Gyumolcsfak virusos, mikoplazmas es rickettsias betegsegei. Budapest: Mezogazdasagi Kiado; 1979.

Nemeth, M.; Kolber, M. Additional evidence on seed transmission of plum pox virus in apricot, peach and plum proved by ELISA Acta Hortic. 130:293-300; 1983.

Nemeth, M. V.; Schmelzer, K. Plum pox virus on Prunus glandulos Thunb. Acta Phytopathol. 7(1/3):193-196; 1972 (In German; English summary).

Pemberton, A. W. Plum pox (sharka disease). Great Britain Ministry Agric. Fish. Food Agric. Dev. Adv. Serv. Leaflet 611, rev. 1980.

Pomazkov, Y.; Abramova, I. Several aspects of epidemiology of sharka virus. Eur. Symp. Fruit Tree Virus Dis. Proc. 8th. Paris: Institut National de la Recherche Agronomique; 1971: 249-253.

Rankovic, M. Reliability of some indicator plants for identification of sharka (plum pox) virus strains. Zast. Bilja 26(133):275-284; 1975 (In Serbo-Croatian; English summary).

- _____. Use of Prunus tomentosa for the detection and differentiation of sharka and other viruses of plum. Acta Phytopathol. Acad. Sci. Hung. 15(1-4):303-308; 1980.
- Roggero, P.; Lenzi, R. Detection of sharka virus on apricot by means of ELISA. Inf. Fitopatol. 35(5):33-35; 1985 (In Italian; English summary).
- Sahtiyanci, S. Sharka virus on plum. FAO Plant Prot. Bull. 17(3):68-69; 1969.
- Sutic, D.; Jordovic, M.; Rankovic, M.; Festic, H. Comparative studies of some sharka (plum pox) virus isolates. Eur. Symp. Fruit Tree Virus Dis. Proc. 8th. Paris: Institut National de la Recherche Agronomique; 1971: 185-192.
- Torrance, L. Use of forced buds to extend the period of serological testing in surveys for fruit tree viruses. Plant Pathol. 30(4):213-216; 1981.
- Trifonov, D. Susceptibility of Prunus insititia to sharka virus and to Polystigma leaf blight. Acta Hortic. 74:229-232; 1978.
- van Oosten, H. J. The isolation of sharka (plum pox) virus from leaves and fruits of plum with herbaceous plants. Neth. J. Plant Pathol. 76(2):99-103; 1970.
- _____. The diagnostic value of inclusion bodies in the fruits of plum trees infected with sharka virus. Eur. Symp. Fruit Tree Virus Dis. Proc. 8th. Paris: Institut National de la Recherche Agronomique; 1971: 211-219.
- _____. Diagnosis of sharka (plum pox) by internal and external fruit symptoms. Neth. J. Plant Pathol. 78(3):99-106; 1972 (Dutch summary).
- Zawadzka, B.; Millikan, D. F. Symptomatology and some biological effects of the sharka disease on common prune in Poland. Phytoprotection 52(2):68-72; 1971.
- Zawadzka, B.; Smolarz, S. Summer hosts of plum aphids as possible host plants of plum pox (sharka) virus. Zesz. Probl. Postepow Nauk. Roln. 214:43-49; 1978 (In Polish; English summary).

